arXiv:1601.07298v1 [math.FA] 27 Jan 2016

# The joint modulus of variation of metric space valued functions and pointwise selection principles

Vyacheslav V. Chistyakov*,[a], Svetlana A. Chistyakova[a]

[a] *Department of Informatics, Mathematics and Computer Science, National Research University Higher School of Economics, Bol'shaya Pechërskaya Street 25/12, Nizhny Novgorod 603155, Russian Federation*

**Abstract**

Given $T \subset \mathbb{R}$ and a metric space $M$, we introduce a nondecreasing sequence of pseudometrics $\{\nu_n\}$ on $M^T$ (the set of all functions from $T$ into $M$), called the joint modulus of variation. We prove that if two sequences of functions $\{f_j\}$ and $\{g_j\}$ from $M^T$ are such that $\{f_j\}$ is pointwise precompact, $\{g_j\}$ is pointwise convergent, and the limit superior of $\nu_n(f_j, g_j)$ as $j \to \infty$ is $o(n)$ as $n \to \infty$, then $\{f_j\}$ admits a pointwise convergent subsequence whose limit is a conditionally regulated function. We illustrate the sharpness of this result by examples (in particular, the assumption on the lim sup is necessary for uniformly convergent sequences $\{f_j\}$ and $\{g_j\}$, and 'almost necessary' when they converge pointwise) and show that most of the known Helly-type pointwise selection theorems are its particular cases.

*Key words:* joint modulus of variation, metric space, regulated function, pointwise convergence, selection principle, generalized variation
*MSC 2000:* 26A45, 28A20, 54C35, 54E50

## 1. Introduction

The purpose of this paper is to present a new sufficient condition (which is almost necessary) on a pointwise precompact sequence $\{f_j\} \equiv \{f_j\}_{j=1}^{\infty}$ of functions $f_j$ mapping a subset $T$ of the real line $\mathbb{R}$ into a metric space $(M, d)$,

*Corresponding author.
*Email addresses:* czeslaw@mail.ru, vchistyakov@hse.ru (Vyacheslav V. Chistyakov), schistyakova@hse.ru (Svetlana A. Chistyakova)

under which the sequence admits a pointwise convergent subsequence. The historically first result in this direction is the classical *Helly Selection Principle*, in which the assumptions are as follows: $T = [a, b]$ is a closed interval, $M = \mathbb{R}$, and $\{f_j\}$ is uniformly bounded and consists of monotone functions ([29], [31, II.8.9–10], [39, VIII.4.2], and [10, Theorem 1.3] if $T \subset \mathbb{R}$ is arbitrary). Since a real function on $T$ of bounded (Jordan) variation is the difference of two nondecreasing bounded functions, Helly's theorem extends to uniformly bounded sequences of functions, whose Jordan's variations are uniformly bounded. Further generalizations of the latter pointwise selection principle are concerned with replacement of Jordan's variation by more general notions of variation ([2, 3], [6]–[10], [15, 16, 22, 25, 26, 32, 33, 38, 41, 43]). In all these papers, the pointwise limit of the extracted subsequence of $\{f_j\}$ is a function of bounded generalized variation (in the corresponding sense), and so, it is a regulated function (with finite one-sided limits at all points of the domain). Note that pointwise selection principles (or the sequential compactness in the topology of pointwise convergence) and regulated functions are of importance in real analysis ([28, 31, 39]), stochastic analysis and generalized integration ([37]), optimization ([1, 36]), set-valued analysis ([2, 10, 19, 20, 30]), and other fields.

A unified approach to the diverse selection principles mentioned above was proposed in [11, 12]. It is based on the notion of *modulus of variation* of a function introduced in [4, 5] (see also [28, 11.3.7]) and does not refer to the uniform boundedness of variations of any kind, and so, can be applied to sequences of non-regulated functions. However, the pointwise limit of the extracted subsequence of $\{f_j\}$ is *again* a regulated function. In order to clarify this situation and expand the amount of sequences having pointwise convergent subsequences, we define the notion of the *joint modulus of variation* for metric space valued functions: this is a certain sequence of pseudometrics $\{\nu_n\}$ on the product set $M^T$ (of all functions from $T$ into $M$). Making use of $\{\nu_n\}$, we obtain a powerful pointwise selection principle (see Theorem 1 in Section 2). Putting $g_j = c$ for all $j \in \mathbb{N}$, where $c : T \to M$ is a constant function, we get the selection principle from [11], which already contains all selection principles alluded to above as particular cases. In contrast to results from [11, 12], the pointwise limit $f$ from Theorem 1 may not be regulated in general—this depends on the limit function $g$, namely, since $\nu_n(f, g) = o(n)$, the function $f$ is only conditionally regulated with respect to $g$ (in short, *$g$-regulated*). In particular, if $g = c$, then $f$ is regulated in the usual sense.

Finally, we point out that following the ideas from [13], Theorem 1 may be extended to sequences of functions with values in a *uniform space* $M$.

The paper is organized as follows. In Section 2, we present necessary definitions and our main result, Theorem 1. In Section 3, we establish essential properties of the joint modulus of variation, which are needed in the proof of Theorem 1 in Section 5. Section 4 is devoted to the study of $g$-regulated (and, in particular, regulated) functions. In the final Section 6, we extend the Helly-type selection theorems from [25] and [22, 32] by exploiting Theorem 1.

## 2. Main result

Let $\varnothing \neq T \subset \mathbb{R}$, $(M, d)$ be a metric space with metric $d$, and $M^T$ denote the set of all functions $f : T \to M$ mapping $T$ into $M$. The letter $c$ stands for a *constant* function $c : T \to M$ (i.e., $c(s) = c(t)$ in $M$ for all $s, t \in T$).

The *joint oscillation* of two functions $f, g \in M^T$ is the quantity

$$|(f, g)(T)| = \sup\big\{|(f, g)(\{s, t\})| : s, t \in T\big\} \in [0, \infty],$$

where

$$|(f, g)(\{s, t\})| = \sup_{z \in M} \big|d(f(s), z) + d(g(t), z) - d(f(t), z) - d(g(s), z)\big| \quad (2.1)$$

is the *joint increment* of $f$ and $g$ on the two-point set $\{s, t\} \subset T$, for which the following two inequalities hold:

$$|(f, g)(\{s, t\})| \leq d(f(s), f(t)) + d(g(s), g(t)), \quad (2.2)$$

$$|(f, g)(\{s, t\})| \leq d(f(s), g(s)) + d(f(t), g(t)). \quad (2.3)$$

Since $|(f, c)(\{s, t\})| = d(f(s), f(t))$ (= the increment of $f$ on $\{s, t\} \subset T$) is independent of $c$, the quantity $|f(T)| = |(f, c)(T)|$ is the usual *oscillation* of $f$ on $T$, also known as the *diameter of the image* $f(T) = \{f(t) : t \in T\} \subset M$. Clearly, by (2.2), $|(f, g)(T)| \leq |f(T)| + |g(T)|$.

We denote by $\mathrm{B}(T; M) = \{f \in M^T : |f(T)| < \infty\}$ the family of all *bounded* functions on $T$ equipped with the *uniform metric* $d_\infty$ given by

$$d_\infty(f, g) = \sup_{t \in T} d(f(t), g(t)) \quad \text{for} \quad f, g \in \mathrm{B}(T; M)$$

($d_\infty$ is an *extended* metric on $M^T$, i.e., may assume the value $\infty$). We have

$$d_\infty(f, g) \leq d(f(s), g(s)) + |f(T)| + |g(T)| \quad \text{for all} \quad s \in T$$

and, by virtue of (2.3), $|(f, g)(T)| \leq 2d_\infty(f, g)$.

If $n \in \mathbb{N}$, we write $\{I_i\}_1^n \prec T$ to denote a collection of $n$ two-point subsets $I_i = \{s_i, t_i\}$ of $T$ $(i = 1, \ldots, n)$ such that $s_1 < t_1 \leq s_2 < t_2 \leq \cdots \leq s_{n-1} < t_{n-1} \leq s_n < t_n$ (so that the intervals $[s_1, t_1], \ldots, [s_n, t_n]$ with end-points in $T$ are non-overlapping). We say that a collection $\{I_i\}_1^n \prec T$ with $I_i = \{s_i, t_i\}$ is a *partition* of $T$ if (setting $t_0 = s_1$) $s_i = t_{i-1}$ for all $i = 1, \ldots, n$, which is written as $\{t_i\}_0^n \prec T$.

The *joint modulus of variation* of two functions $f, g \in M^T$ is the sequence $\{\nu_n(f,g)\}_{n=1}^{\infty} \subset [0, \infty]$ defined by

$$\nu_n(f,g) = \sup\left\{\sum_{i=1}^{n} |(f,g)(I_i)| : \{I_i\}_1^n \prec T\right\} \quad \text{for all} \quad n \in \mathbb{N}, \tag{2.4}$$

where $|(f,g)(I_i)| = |(f,g)(\{s_i, t_i\})|$ is the quantity from (2.1) if $I_i = \{s_i, t_i\}$ (for finite $T$ with the number of elements $\#(T) \geq 2$, we make use of (2.4) for $n \leq \#(T) - 1$, and set $\nu_n(f,g) = \nu_{\#(T)-1}(f,g)$ for all $n > \#(T) - 1$).

Note that, given $f, g \in M^T$, we have $\nu_1(f,g) = |(f,g)(T)|$ and

$$\nu_1(f,g) \leq \nu_n(f,g) \leq n\nu_1(f,g) \quad \text{for all} \quad n \in \mathbb{N}. \tag{2.5}$$

Further properties of the joint modulus of variation are presented in Section 3.

For a sequence of functions $\{f_j\} \subset M^T$ and $f \in M^T$, we write: (a) $f_j \to f$ on $T$ to denote the *pointwise* (or *everywhere*) *convergence* of $\{f_j\}$ to $f$ (that is, $\lim_{j\to\infty} d(f_j(t), f(t)) = 0$ for all $t \in T$); (b) $f_j \rightrightarrows f$ on $T$ to denote the *uniform convergence* of $\{f_j\}$ to $f$ meaning, as usual, that $\lim_{j\to\infty} d_\infty(f_j, f) = 0$. The uniform convergence implies the pointwise convergence, but not vice versa. Recall that a sequence $\{f_j\} \subset M^T$ is said to be *pointwise precompact* on $T$ if the closure in $M$ of the set $\{f_j(t) : j \in \mathbb{N}\}$ is compact for all $t \in T$.

Making use of E. Landau's notation, given a sequence, $\{\mu_n\}_{n=1}^{\infty} \subset \mathbb{R}$, we write $\mu_n = o(n)$ to denote the condition $\lim_{n\to\infty} \mu_n/n = 0$.

Our main result, a *pointwise selection principle* for metric space valued functions in terms of the joint modulus of variation, is as follows.

**Theorem 1.** *Let $\varnothing \neq T \subset \mathbb{R}$ and $(M,d)$ be a metric space. Suppose $\{f_j\}, \{g_j\} \subset M^T$ are two sequences of functions such that*

(a) *$\{f_j\}$ is pointwise precompact on $T$,*

(b) *$\{g_j\}$ is pointwise convergent on $T$ to a function $g \in M^T$,*

*and*

$$\mu_n \equiv \limsup_{j\to\infty} \nu_n(f_j, g_j) = o(n). \tag{2.6}$$

*Then, there is a subsequence of $\{f_j\}$, which converges pointwise on $T$ to a function $f \in M^T$ such that $\nu_n(f,g) \le \mu_n$ for all $n \in \mathbb{N}$.*

This theorem will be proved in Section 5. Now, a few remarks are in order. Given $f \in M^T$ and a constant function $c : T \to M$, the quantity

$$\nu_n(f) \equiv \nu_n(f,c) = \sup\left\{\sum_{i=1}^{n} d(f(s_i), f(t_i)) : \{I_i\}_1^n \prec T\right\} \tag{2.7}$$

(with $I_i = \{s_i, t_i\}$) is independent of $c$, and the sequence $\{\nu_n(f)\}_{n=1}^{\infty} \subset [0,\infty]$ is known as the *modulus of variation* of $f$ in the sense of Chanturiya ([4, 5, 11, 12, 13, 28]). It characterizes regulated (or proper) functions on $T = [a,b]$ as follows. We say that $f : [a,b] \to M$ is *regulated* and write $f \in \mathrm{Reg}([a,b]; M)$ if $d(f(s), f(t)) \to 0$ as $s, t \to \tau - 0$ for every $a < \tau \le b$, and $d(f(s), f(t)) \to 0$ as $s, t \to \tau' + 0$ for every $a \le \tau' < b$ (and so, by Cauchy's criterion, one-sided limits $f(\tau - 0), f(\tau' + 0) \in M$ exist provided $M$ is complete). We have

$$\mathrm{Reg}([a,b]; M) = \{f \in M^{[a,b]} : \nu_n(f) = o(n)\} \tag{2.8}$$

(more general characterizations for dense subsets $T$ of $[a,b]$ can be found in [12, 13]). A certain relationship between characterizations of regulated functions and pointwise selection principles is exhibited in [18].

## 3. The joint modulus of variation

We begin by studying the joint increment (2.1), whose properties are gathered in the following lemma.

**Lemma 1.** *Given $f, g, h \in M^T$ and $s, t \in T$, we have*:

(a) $|(f,f)(\{s,t\})| = 0$;
(b) $|(f,g)(\{s,t\})| = |(g,f)(\{s,t\})|$;
(c) $|(f,g)(\{s,t\})| \le |(f,h)(\{s,t\})| + |(h,g)(\{s,t\})|$;
(d) $d(f(s), f(t)) \le d(g(s), g(t)) + |(f,g)(\{s,t\})|$;
(e) $d(f(t), g(t)) \le d(f(s), g(s)) + |(f,g)(\{s,t\})|$.

**Proof**. Properties (a), (b), and (c), showing that $(f,g) \mapsto |(f,g)(\{s,t\})|$ is a *pseudometric* on $M^T$, are straightforward. To establish (d) and (e), take into account equality $d(x,y) = \max_{z \in M} |d(x,z) - d(y,z)|$. □

**Remark 1.** (a) If $|(f,g)(\{s,t\})| = 0$, then (d), (e), and (b) imply equalities $d(f(s),f(t)) = d(g(s),g(t))$ and $d(f(t),g(t)) = d(f(s),g(s))$. In addition to Lemma 1, the function $(s,t) \mapsto |(f,g)(\{s,t\})|$ is a pseudometric on $T$.

(b) If $F(z)$ denotes the absolute value under the supremum sign in (2.1), then $F : M \to [0,\infty)$ and $|F(z) - F(z_0)| \le 4d(z,z_0)$ for all $z, z_0 \in M$.

(c) By Lemma 1(d), $|f(T)| \le |g(T)| + |(f,g)(T)| = |g(T)| + \nu_1(f,g)$. So,

$$\big||f(T)| - |g(T)|\big| \le |(f,g)(T)| \le |f(T)| + |g(T)|, \qquad f,g \in \mathrm{B}(T;M).$$

Moreover, it follows from Lemma 1(e) that

$$d_\infty(f,g) \le d(f(s),g(s)) + |(f,g)(T)| \le 3d_\infty(f,g) \quad \text{for all} \quad s \in T.$$

(d) Suppose the triple $(M,d,+)$ is a *metric semigroup* ([10, Section 4]), i.e., $(M,d)$ is a metric space, $(M,+)$ is an Abelian semigroup with the operation of addition $+$, and $d(x,y) = d(x+z,y+z)$ for all $x,y,z \in M$. Then, the joint increment (2.1) may be alternatively replaced by

$$|(f,g)(\{s,t\})| = d(f(s)+g(t), f(t)+g(s)). \tag{3.1}$$

The joint modulus of variation (2.4) involving (3.1) was employed in [17]. Furthermore, if $(M, \|\cdot\|)$ is a normed linear space (over $\mathbb{R}$ or $\mathbb{C}$), we may set

$$|(f,g)(\{s,t\})| = \|f(s)+g(t)-f(t)-g(s)\| = \|(f-g)(s)-(f-g)(t)\|. \tag{3.2}$$

Quantities (3.1) and (3.2) have the same properties as (2.1): see (2.2), (2.3), Lemma 1 and Remark 1(a). In the sequel, we make use of more general quantity (2.1).

If $f,g \in M^T$, $n \in \mathbb{N}$ and $\varnothing \ne E \subset T$, we set $\nu_n(f,g;E) = \nu_n(f|_E, g|_E)$, where $f|_E \in M^E$ is the restriction of $f$ to $E$ (i.e., $f|_E(t) = f(t)$ for all $t \in E$). Accordingly, $\nu_n(f,g) = \nu_n(f,g;T)$.

The following properties of the joint modulus of variation are immediate. The sequence $\{\nu_n(f,g)\}_{n=1}^\infty$ is nondecreasing, $\nu_{n+m}(f,g) \le \nu_n(f,g) + \nu_m(f,g)$ for all $n,m \in \mathbb{N}$, and $\nu_n(f,g;E) \le \nu_n(f,g;T)$ provided $n \in \mathbb{N}$ and $E \subset T$. It follows from (2.4) and Lemma 1(a)–(c) that, for every $n \in \mathbb{N}$, the function $(f,g) \mapsto \nu_n(f,g)$ is a *pseudometric* on $M^T$ (possibly assuming infinite values) and, in particular (cf. (2.4) and (2.7)),

$$\nu_n(f,g) \le \nu_n(f) + \nu_n(g) \quad \text{and} \quad \nu_n(f) \le \nu_n(g) + \nu_n(f,g) \tag{3.3}$$

for all $n \in \mathbb{N}$ and $f,g \in M^T$. Furthemore, if $f,g \in \mathrm{B}(T;M)$, then, by (2.5), the sequence $\{\nu_n(f,g)/n\}_{n=1}^\infty$ is bounded in $[0,\infty)$.

Essential properties of the joint modulus of variation are presented in

**Lemma 2.** *Given $n \in \mathbb{N}$, $f, g \in M^T$, and $\varnothing \neq E \subset T$, we have*:

(a) $|(f,g)(\{s,t\})| + \nu_n(f,g;E_s^-) \leq \nu_{n+1}(f,g;E_t^-)$ *for all* $s,t \in E$ *with* $s \leq t$, *where* $E_\tau^- = (-\infty,\tau] \cap E$ *for* $\tau \in E$;

(b) $\nu_{n+1}(f,g;E) \leq \nu_n(f,g;E) + \dfrac{\nu_{n+1}(f,g;E)}{n+1}$;

(c) *if* $\{f_j\}, \{g_j\} \subset M^T$ *are such that* $f_j \to f$ *and* $g_j \to g$ *on* $E$, *then* $\nu_n(f,g;E) \leq \liminf_{j\to\infty} \nu_n(f_j,g_j;E)$;

(d) *if* $\{f_j\}, \{g_j\} \subset M^T$ *are such that* $f_j \rightrightarrows f$ *and* $g_j \rightrightarrows g$ *on* $E$, *then* $\nu_n(f,g;E) = \lim_{j\to\infty} \nu_n(f_j,g_j;E)$.

**Proof**. (a) We may assume that $s < t$. Let $\{I_i\}_1^n \prec E_s^-$. Setting $I_0 = \{s,t\}$, we find $\{I_i\}_0^n \prec E_t^-$, and so,

$$|(f,g)(I_0)| + \sum_{i=1}^{n} |(f,g)(I_i)| \leq \nu_{n+1}(f,g;E_t^-).$$

The inequality in (a) follows by taking the supremum over all $\{I_i\}_1^n \prec E_s^-$.

(b) We may assume that $\nu_{n+1}(f,g;E)$ is finite, and apply the idea from [5, Lemma]. Given $\varepsilon > 0$, there is $\{I_i\}_1^{n+1} \prec E$ (depending on $\varepsilon$) such that

$$\sum_{i=1}^{n+1} |(f,g)(I_i)| \leq \nu_{n+1}(f,g;E) \leq \sum_{i=1}^{n+1} |(f,g)(I_i)| + \varepsilon.$$

Setting $a_0 = \min_{1\leq i\leq n+1} |(f,g)(I_i)|$, the left-hand side inequality implies $(n+1)a_0 \leq \nu_{n+1}(f,g;E)$. The right-hand side inequality gives

$$\nu_{n+1}(f,g;E) \leq \nu_n(f,g;E) + a_0 + \varepsilon,$$

from which our inequality follows due to the arbitrariness of $\varepsilon > 0$.

(c) First, we note that, given $j \in \mathbb{N}$ and $s,t \in T$, we have

$$\big||(f_j,g_j)(\{s,t\})| - |(f,g)(\{s,t\})|\big| \leq d(f_j(s),f(s)) + d(f_j(t),f(t)) + d(g_j(s),g(s)) + d(g_j(t),g(t)). \quad (3.4)$$

In fact, Lemma 1(c) and inequality (2.3) imply

$$\begin{aligned}|(f_j,g_j)(\{s,t\})| &\le |(f_j,f)(\{s,t\})| + |(f,g)(\{s,t\})| + |(g,g_j)(\{s,t\})| \\ &\le d(f_j(s),f(s)) + d(f_j(t),f(t)) + |(f,g)(\{s,t\})| \\ &\quad + d(g(s),g_j(s)) + d(g(t),g_j(t)). \end{aligned} \tag{3.5}$$

Exchanging $f_j$ and $f$ as well as $g_j$ and $g$, we obtain (3.4).

From the pointwise convergence of $\{f_j\}$ and $\{g_j\}$ and (3.4), we find

$$\lim_{j\to\infty} |(f_j,g_j)(\{s,t\})| = |(f,g)(\{s,t\})| \quad \text{for all} \quad s,t\in E.$$

By definition (2.4), given $\{I_i\}_1^n \prec E$, we have

$$\sum_{i=1}^n |(f_j,g_j)(I_i)| \le \nu_n(f_j,g_j;E) \quad \text{for all} \quad j\in\mathbb{N}.$$

Passing to the limit inferior as $j\to\infty$, we get

$$\sum_{i=1}^n |(f,g)(I_i)| \le \liminf_{j\to\infty} \nu_n(f_j,g_j;E). \tag{3.6}$$

Since $\{I_i\}_1^n \prec E$ is arbitrary, it remains to take into account (2.4).

(d) It follows from (3.5) that, for any $s,t\in E$ and $j\in\mathbb{N}$,

$$|(f_j,g_j)(\{s,t\})| \le 2\sup_{\tau\in E} d(f_j(\tau),f(\tau)) + |(f,g)(\{s,t\}) + 2\sup_{\tau\in E} d(g_j(\tau),g(\tau)),$$

and so, definition (2.4) implies

$$\nu_n(f_j,g_j;E) \le 2n\sup_{\tau\in E} d(f_j(\tau),f(\tau)) + \nu_n(f,g;E) + 2n\sup_{\tau\in E} d(g_j(\tau),g(\tau)) \tag{3.7}$$

for all $j\in\mathbb{N}$. Passing to the limit superior as $j\to\infty$, we get

$$\limsup_{j\to\infty} \nu_n(f_j,g_j;E) \le \nu_n(f,g;E).$$

Now, the equality in (d) is a consequence of Lemma 2(c). □

**Remark 2.** If the value $\nu_1(f,g;E) = |(f,g)(E)|$ (see (2.5)) is finite for an $E\subset T$ (e.g., when $f,g\in \mathrm{B}(E;M)$), inequality in Lemma 2(b) is equivalent to

$$\frac{\nu_{n+1}(f,g;E)}{n+1} \le \frac{\nu_n(f,g;E)}{n}.$$

Thus, the limit $\lim_{n\to\infty} \nu_n(f,g;E)/n$ always exists in $[0,\infty)$.

## 4. Conditionally regulated functions

Since $\nu_n = \nu_n(\cdot,\cdot)$ is a (extended) pseudometric on $M^T$, we may introduce an equivalence relation $\sim$ on $M^T$ as follows: given $f, g \in M^T$, we set

$$f \sim g \quad \text{if and only if} \quad \nu_n(f,g) = o(n).$$

The equivalence class $\mathrm{R}(g) = \{f \in M^T : f \sim g\}$ of a function $g \in M^T$ is called the *regularity class* of $g$, and any representative $f \in \mathrm{R}(g)$ is called a *conditionally regulated* or, more precisely, *$g$-regulated* function. This terminology is justified by (2.8): in the framework of the product set $M^{[a,b]}$, we have $\mathrm{Reg}([a,b]; M) = \mathrm{R}(c)$ for any constant function $c : [a,b] \to M$.

Note that, in Theorem 1, condition '$\nu_n(f,g) \le \mu_n$ for all $n \in \mathbb{N}$' means that $f \in \mathrm{R}(g)$, and so, the class $\mathrm{R}(g)$ is worth studying in more detail.

**Theorem 2.** *Given a function $g \in M^T$, we have*:

(a) *$g \in \mathrm{B}(T; M)$ if and only if $\mathrm{R}(g) \subset \mathrm{B}(T; M)$*;

(b) *$\mathrm{R}(g)$ is closed with respect to the uniform convergence, but not closed with respect to the pointwise convergence in general*;

(c) *if $(M, d)$ is a complete metric space, then the pair $(\mathrm{R}(g), d_\infty)$ is also a complete metric space.*

**Proof**. (a) The sufficiency is clear, because $g \in \mathrm{R}(g)$. Now, suppose that $g \in \mathrm{B}(T; M)$, so that, by (2.7), $\nu_1(g) = |(g, c)(T)| = |g(T)| < \infty$. Given $f \in \mathrm{R}(g)$, $\nu_n(f,g) = o(n)$, and so, $\nu_{n_0}(f,g) \le n_0$ for some $n_0 \in \mathbb{N}$. It follows from (3.3) and (2.5) that

$$|f(T)| = \nu_1(f) \le \nu_1(g) + \nu_1(f,g) \le |g(T)| + \nu_{n_0}(f,g) \le |g(T)| + n_0 < \infty,$$

which implies $f \in \mathrm{B}(T; M)$.

(b) It is to be shown that if $\{f_j\} \subset \mathrm{R}(g)$ and $f_j \rightrightarrows f$ on $T$ with $f \in M^T$, then $f \in \mathrm{R}(g)$. We will prove a little bit more: namely, if $\{f_j\}, \{g_j\} \subset M^T$ are such that $f_j \in \mathrm{R}(g_j)$ for all $j \in \mathbb{N}$, $f_j \rightrightarrows f$ and $g_j \rightrightarrows g$ on $T$ with $f, g \in M^T$, then $f \in \mathrm{R}(g)$ (the previous assertion follows if $g_j = g$ for all $j \in \mathbb{N}$). In fact, exchanging $f_j$ and $f$, and $g_j$ and $g$ in (3.7), we get

$$\frac{\nu_n(f,g)}{n} \le 2d_\infty(f, f_j) + \frac{\nu_n(f_j, g_j)}{n} + 2d_\infty(g, g_j), \qquad n, j \in \mathbb{N}.$$

By the uniform convergence of $\{f_j\}$ and $\{g_j\}$, given $\varepsilon > 0$, there is a number $j_0 = j_0(\varepsilon) \in \mathbb{N}$ such that $d_\infty(f, f_{j_0}) \le \varepsilon$ and $d_\infty(g, g_{j_0}) \le \varepsilon$. Since $f_{j_0}$ is in $\mathrm{R}(g_{j_0})$, we have $\nu_n(f_{j_0}, g_{j_0}) = o(n)$, and so, there exists $n_0 = n_0(\varepsilon) \in \mathbb{N}$ such that $\nu_n(f_{j_0}, g_{j_0})/n \le \varepsilon$ for all $n \ge n_0$. The estimate above with $j = j_0$ implies $\nu_n(f, g)/n \le 5\varepsilon$, $n \ge n_0$, which means that $\nu_n(f, g) = o(n)$ and $f \in \mathrm{R}(g)$.

As for the pointwise convergence, consider a sequence of real step functions converging pointwise to the Dirichlet function (=the characteristic function of the rationals $\mathbb{Q}$) on $T = [0, 1]$ (see [11, Examples 4, 5] and Example 2(a) in Section 5).

(c) First, we show that $d_\infty(f, f') < \infty$ for all $f, f' \in \mathrm{R}(g)$. In fact, since $f \sim f'$, we have $\nu_n(f, f') = o(n)$, and so, $\nu_{n_0}(f, f') \le n_0$ for some $n_0 \in \mathbb{N}$. Given $s \in T$, it follows from Remark 1(c) and (2.5) that

$$d_\infty(f, f') \le d(f(s), f'(s)) + \nu_1(f, f') \le d(f(s), f'(s)) + \nu_{n_0}(f, f') < \infty.$$

The metric axioms for $d_\infty$ on $\mathrm{R}(g)$ are verified in a standard way.

In order to prove that $\mathrm{R}(g)$ is complete, suppose $\{f_j\} \subset \mathrm{R}(g)$ is a Cauchy sequence, i.e., $d_\infty(f_j, f_k) \to 0$ as $j, k \to \infty$. Since $d(f_j(t), f_k(t)) \le d_\infty(f_j, f_k)$ for all $t \in T$ and $(M, d)$ is complete, there exists $f \in M^T$ such that $f_j \to f$ on $T$. Noting that $f_j \to f_j$ and $f_k \to f$ on $T$ as $k \to \infty$ (and arguing as in (3.6)), we get

$$d_\infty(f_j, f) \le \liminf_{k\to\infty} d_\infty(f_j, f_k) = \lim_{k\to\infty} d_\infty(f_j, f_k) < \infty \quad \text{for all} \quad j \in \mathbb{N}.$$

Since the sequence $\{f_j\}$ is $d_\infty$-Cauchy, we find

$$\limsup_{j\to\infty} d_\infty(f_j, f) \le \lim_{j\to\infty} \lim_{k\to\infty} d_\infty(f_j, f_k) = 0.$$

Thus, $\lim_{j\to\infty} d_\infty(f_j, f) = 0$, and so, $f_j \rightrightarrows f$ on $T$. Applying item (b) of this Theorem, we conclude that $f \in \mathrm{R}(g)$. □

A traditionally important class of regulated functions is the space of functions of bounded Jordan variation, $\mathrm{BV}(T; M)$, which is introduced by means of the joint modulus of variation as follows.

Since the sequence $\{\nu_n(f, g)\}_{n=1}^\infty$ is nondecreasing for all $f, g \in M^T$, the quantity (finite or not) $V(f, g) = \lim_{n\to\infty} \nu_n(f, g) = \sup_{n\in\mathbb{N}} \nu_n(f, g)$ is called the *joint variation* of functions $f$ and $g$ on $T$. The value $V(f) \equiv V(f, c)$

is independent of a constant function $c : T \to M$ and is the usual Jordan *variation* of $f$ on $T$:

$$V(f) = \sup\Big\{\sum_{i=1}^{n} d(f(t_i), f(t_{i-1})) : n \in \mathbb{N} \text{ and } \{t_i\}_0^n \prec T\Big\},$$

the supremum being taken over all partitions $\{t_i\}_0^n$ of $T$ (cf. Section 2). The set $\mathrm{BV}(T; M) = \{f \in M^T : V(f) < \infty\}$ is contained in $\mathrm{B}(T; M) \cap \mathrm{R}(c)$ (in fact, $|f(T)| = \nu_1(f) \le V(f)$ and $\nu_n(f, c)/n \le V(f)/n$ for all $f \in \mathrm{BV}(T; M)$).

The following notion of *$\varepsilon$-variation* $V_\varepsilon(f)$, due to Fraňková [25, Section 3], provides an alternative characterization (cf. (2.8)) of regulated functions: given $f \in M^T$ and $\varepsilon > 0$, set

$$V_\varepsilon(f) = \inf\big\{V(g) : g \in \mathrm{BV}(T; M) \text{ and } d_\infty(f, g) \le \varepsilon\big\} \quad (\inf \varnothing = \infty). \tag{4.1}$$

It was shown in [25, Proposition 3.4] (for $T = [a, b]$) that

$$\mathrm{Reg}([a, b]; M) = \{f \in M^{[a,b]} : V_\varepsilon(f) < \infty \text{ for all } \varepsilon > 0\} \tag{4.2}$$

(although it was assumed in [25] that $M = \mathbb{R}^N$, the proof of the last assertion carries over to any metric space $M$, cf. [11, Lemma 3]).

The notion of $\varepsilon$-variation will be needed in Section 6.

**Example 1.** Given $x, y \in M$ with $x \ne y$, let $f = \mathcal{D}_{x,y} : T = [0, 1] \to M$ be the Dirichlet-type function of the form:

$$\mathcal{D}_{x,y}(t) = \begin{cases} x & \text{if } \ t \in [0, 1] \text{ is rational,} \\ y & \text{if } \ t \in [0, 1] \text{ is irrational.} \end{cases} \tag{4.3}$$

Clearly, $f \notin \mathrm{Reg}([0, 1]; M)$. Moreover (cf. (4.2)), we have

$$V_\varepsilon(f) = \infty \text{ if } 0 < \varepsilon < d(x, y)/2, \quad \text{and} \quad V_\varepsilon(f) = 0 \text{ if } \varepsilon \ge d(x, y). \tag{4.4}$$

To see this, first note that, given $g \in M^{[0,1]}$, inequality $d_\infty(f, g) \le \varepsilon$ is equivalent to the following two conditions:

$$d(x, g(s)) \le \varepsilon \ \ \forall\, s \in [0, 1] \cap \mathbb{Q}, \quad \text{and} \quad d(y, g(t)) \le \varepsilon \ \ \forall\, t \in [0, 1] \setminus \mathbb{Q}. \tag{4.5}$$

Suppose $0 < \varepsilon < d(x, y)/2$. To show that $d_\infty(f, g) \le \varepsilon$ implies $V(g) = \infty$, we let $n \in \mathbb{N}$, and $\{t_i\}_0^{2n} \prec [0, 1]$ be a partition of $[0, 1]$ such that points

$\{t_{2i}\}_{i=0}^n$ are rational and points $\{t_{2i-1}\}_{i=1}^n$ are irrational. By the triangle inequality for $d$ and (4.5), we get

$$\begin{aligned} V(g) &\geq \sum_{i=1}^{2n} d(g(t_i), g(t_{i-1})) \geq \sum_{i=1}^{n} d(g(t_{2i}), g(t_{2i-1})) \\ &\geq \sum_{i=1}^{n} \big(d(x,y) - d(x, g(t_{2i})) - d(g(t_{2i-1}), y)\big) \geq n\big(d(x,y) - 2\varepsilon\big). \end{aligned}$$

If $\varepsilon \geq d(x,y)$, we set $g(t) = x$ (or $g(t) = y$) for all $t \in [0,1]$, so that (4.5) is satisfied and $V(g) = 0$. Thus, $V_\varepsilon(f) = 0$.

The second assertion in (4.4) can be refined, provided

$$d(x,y)/2 = \max\{d(x,z_0), d(y,z_0)\} \quad \text{for some} \quad z_0 \in M. \tag{4.6}$$

In fact, we may set $g(t) = z_0$ for all $t \in [0,1]$, so that (4.5) holds whenever $d(x,y)/2 \leq \varepsilon$, and $V(g) = 0$. This implies $V_\varepsilon(f) = 0$ for all $\varepsilon \geq d(x,y)/2$.

A few remarks concerning condition (4.6) are in order. Since

$$d(x,y) \leq d(x,z) + d(z,y) \leq 2\max\{d(x,z), d(y,z)\} \quad \text{for all} \quad z \in M,$$

condition (4.6) refers to a certain form of 'convexity' of $M$ (which is not restrictive for our purposes). For instance, if $(M, \|\cdot\|)$ is a normed linear space with $d(x,y) = \|x - y\|$, we may set $z_0 = (x+y)/2$. More generally, by Menger's Theorem ([35], [27, Example 2.7]), if a metric space $(M,d)$ is complete and *metrically convex* (i.e., given $x, y \in M$ with $x \neq y$, there is $z \in M$ such that $x \neq z \neq y$ and $d(x,y) = d(x,z) + d(z,y)$), then, for any $x, y \in M$, there is an isometry $\varphi : [0, d(x,y)] \to M$ such that $\varphi(0) = x$ and $\varphi(d(x,y)) = y$. In this case, we set $z_0 = \varphi(d(x,y)/2)$. More examples of metrically convex metric spaces can be found in [21, 24].

Finally, if $M = \{x, y\}$, then condition (4.6) is not satisfied, and we have $V_\varepsilon(f) = \infty$ for all $0 < \varepsilon < d(x,y)$, which is a consequence of (4.5).

## 5. Proof of the main result

**Proof of Theorem 1**. With no loss of generality we may assume that $T$ is uncountable; otherwise, by virtue of assumption (a) and the standard Cantor diagonal procedure, we extract a pointwise convergent subsequence of $\{f_j\}$ and apply Lemma 2(c). Note that $\mu_n$ is finite for all $n \in \mathbb{N}$: in fact, $\mu_n \leq n$ whenever $n \geq n_0$ for some $n_0 \in \mathbb{N}$ and, since $n \mapsto \nu_n(f_j, g_j)$ is nondecreasing for all $j \in \mathbb{N}$, we have $\mu_n \leq \mu_{n_0} \leq n_0$ for all $1 \leq n < n_0$.

We divide the rest of the proof into four steps for clarity.

*Step 1.* Let us show that there is a subsequence of $\{j\}_{j=1}^{\infty}$, again denoted by $\{j\}$, and a nondecreasing sequence $\{\alpha_n\}_{n=1}^{\infty} \subset [0,\infty)$ such that

$$\lim_{j\to\infty} \nu_n(f_j, g_j) = \alpha_n \le \mu_n \quad \text{for all} \quad n \in \mathbb{N}. \tag{5.1}$$

We set $\alpha_1 = \mu_1$. The definition (2.6) of $\mu_1$ implies that there is an increasing sequence $\{J_1(j)\}_{j=1}^{\infty} \subset \mathbb{N}$ (i.e., a subsequence of $\{j\}_{j=1}^{\infty}$) such that $\nu_1(f_{J_1(j)}, g_{J_1(j)}) \to \alpha_1$ as $j \to \infty$. Setting $\alpha_2 = \limsup_{j\to\infty} \nu_2(f_{J_1(j)}, g_{J_1(j)})$, we find $\alpha_2 \le \mu_2$, and there is a subsequence $\{J_2(j)\}_{j=1}^{\infty}$ of $\{J_1(j)\}_{j=1}^{\infty}$ such that $\nu_2(f_{J_2(j)}, g_{J_2(j)}) \to \alpha_2$ as $j \to \infty$. Inductively, if $n \ge 3$ and a subsequence $\{J_{n-1}(j)\}_{j=1}^{\infty}$ of $\{j\}_{j=1}^{\infty}$ is already chosen, we define $\alpha_n$ as the limit superior of $\nu_n(f_{J_{n-1}(j)}, g_{J_{n-1}(j)})$ as $j \to \infty$, so that $\alpha_n \le \mu_n$. Now, we pick a subsequence $\{J_n(j)\}_{j=1}^{\infty}$ of $\{J_{n-1}(j)\}_{j=1}^{\infty}$ such that $\nu_n(f_{J_n(j)}, g_{J_n(j)}) \to \alpha_n$ as $j \to \infty$. Noting that the sequence $\{J_j(j)\}_{j=n}^{\infty}$ is a subsequence of $\{J_n(j)\}_{j=1}^{\infty}$ (for all $n \in \mathbb{N}$) and denoting the diagonal sequences $\{f_{J_j(j)}\}_{j=1}^{\infty}$ and $\{g_{J_j(j)}\}_{j=1}^{\infty}$ again by $\{f_j\}$ and $\{g_j\}$, respectively, we obtain (5.1).

In the sequel, the set of all nondecreasing bounded functions mapping $T$ into $\mathbb{R}^+ = [0,\infty)$ is denoted by $\mathrm{Mon}(T;\mathbb{R}^+)$.

*Step 2.* In this step, we prove that there are subsequences of $\{f_j\}$ and $\{g_j\}$ from (5.1), again denoted by $\{f_j\}$ and $\{g_j\}$, respectively, and a sequence of functions $\{\beta_n\}_{n=1}^{\infty} \subset \mathrm{Mon}(T;\mathbb{R}^+)$ such that

$$\lim_{j\to\infty} \nu_n(f_j, g_j; T_t^-) = \beta_n(t) \quad \text{for all } n \in \mathbb{N} \text{ and } t \in T, \tag{5.2}$$

where $T_t^- = \{s \in T : s \le t\}$ for $t \in T$.

Note that, for each $n \in \mathbb{N}$, the function $t \mapsto \nu_n(f_j, g_j; T_t^-)$ is nondecreasing on $T$, and $\nu_n(f_j, g_j; T_t^-) \le \nu_n(f_j, g_j)$ for all $t \in T$ and $n \in \mathbb{N}$. By virtue of (5.1), there is a sequence $\{C_n\}_{n=1}^{\infty} \subset \mathbb{R}^+$ such that $\nu_n(f_j, g_j) \le C_n$ for all $n, j \in \mathbb{N}$. In what follows, we apply the diagonal procedure once again.

The sequence $\{t \mapsto \nu_1(f_j, g_j; T_t^-)\}_{j=1}^{\infty} \subset \mathrm{Mon}(T;\mathbb{R}^+)$ is uniformly bounded by constant $C_1$, and so, by Helly's Selection Principle, there are an increasing sequence $\{K_1(j)\}_{j=1}^{\infty} \subset \mathbb{N}$ (i.e., a subsequence of $\{j\}_{j=1}^{\infty}$) and a function $\beta_1 \in \mathrm{Mon}(T;\mathbb{R}^+)$ such that $\nu_1(f_{K_1(j)}, g_{K_1(j)}; T_t^-) \to \beta_1(t)$ as $j \to \infty$ for all $t \in T$. The sequence $\{t \mapsto \nu_2(f_{K_1(j)}, g_{K_1(j)}; T_t^-)\}_{j=1}^{\infty} \subset \mathrm{Mon}(T;\mathbb{R}^+)$ is uniformly bounded on $T$ by constant $C_2$, and so, again by Helly's Theorem, there are a subsequence $\{K_2(j)\}_{j=1}^{\infty}$ of $\{K_1(j)\}_{j=1}^{\infty}$ and a function $\beta_2 \in \mathrm{Mon}(T;\mathbb{R}^+)$

such that $\nu_2(f_{K_2(j)}, g_{K_2(j)}; T_t^-) \to \beta_2(t)$ as $j \to \infty$ for all $t \in T$. Inductively, if $n \geq 3$ and a subsequence $\{K_{n-1}(j)\}_{j=1}^\infty$ of $\{j\}_{j=1}^\infty$ and a function $\beta_{n-1} \in \mathrm{Mon}(T; \mathbb{R}^+)$ are already chosen, we apply the Helly Theorem to the sequence of functions $\{t \mapsto \nu_n(f_{K_{n-1}(j)}, g_{K_{n-1}(j)}; T_t^-)\}_{j=1}^\infty \subset \mathrm{Mon}(T; \mathbb{R}^+)$, which is uniformly bounded on $T$ by constant $C_n$: there are a subsequence $\{K_n(j)\}_{j=1}^\infty$ of $\{K_{n-1}(j)\}_{j=1}^\infty$ and a function $\beta_n \in \mathrm{Mon}(T; \mathbb{R}^+)$ such that $\nu_n(f_{K_n(j)}, g_{K_n(j)}; T_t^-) \to \beta_n(t)$ as $j \to \infty$ for all $t \in T$. Since $\{K_j(j)\}_{j=n}^\infty$ is a subsequence of $\{K_n(j)\}_{j=1}^\infty$ (for all $n \in \mathbb{N}$), the diagonal sequences $\{f_{K_j(j)}\}_{j=1}^\infty$ and $\{g_{K_j(j)}\}_{j=1}^\infty$, again denoted by $\{f_j\}$ and $\{g_j\}$, respectively, satisfy condition (5.2).

*Step 3.* Let $Q$ be an at most countable dense subset of $T$. Note that $Q$ contains all isolated (= non-limit) points of $T$ (i.e., points $t \in T$ such that the intervals $(t-\delta, t)$ and $(t, t+\delta)$ lie in $\mathbb{R} \setminus T$ for some $\delta > 0$). The set $Q_n \subset T$ of discontinuity points of nondecreasing function $\beta_n$ is at most countable. Setting $S = Q \cup \bigcup_{n=1}^\infty Q_n$, we find that $S$ is an at most countable dense subset of $T$ and

$$\beta_n \text{ is continuous at all points of } T \setminus S \text{ for all } n \in \mathbb{N}. \tag{5.3}$$

Since the set $\{f_j(t) : j \in \mathbb{N}\}$ is precompact in $M$ for all $t \in T$, and $S \subset T$ is at most countable, we may assume (applying the diagonal procedure again and passing to a subsequence of $\{f_j\}$ if necessary) that, given $s \in S$, there is a point $f(s) \in M$ such that $d(f_j(s), f(s)) \to 0$ as $j \to \infty$. In this way, we obtain a function $f : S \to M$.

*Step 4.* Now, we show that, for every $t \in T \setminus S$, the sequence $\{f_j(t)\}_{j=1}^\infty$ converges in $M$. For this, we prove that this sequence is Cauchy in $M$, i.e., $d(f_j(t), f_k(t)) \to 0$ as $j, k \to \infty$. Let $\varepsilon > 0$ be arbitrarily fixed. By assumption (2.6), $\mu_n/n \to 0$ as $n \to \infty$, so we choose and fix a number $n = n(\varepsilon) \in \mathbb{N}$ such that

$$\frac{\mu_{n+1}}{n+1} \leq \varepsilon.$$

By property (5.1), there is a number $j_1 = j_1(\varepsilon, n) \in \mathbb{N}$ such that

$$\nu_{n+1}(f_j, g_j) \leq \alpha_{n+1} + \varepsilon \leq \mu_{n+1} + \varepsilon \quad \text{for all} \quad j \geq j_1.$$

The definition of the set $S$ and (5.3) imply that the point $t$ is a limit point for $T$ and, at the same time, a point of continuity of the function $\beta_n$. By the density of $S$ in $T$, there is $s = s(\varepsilon, n, t) \in S$ such that

$$|\beta_n(t) - \beta_n(s)| \leq \varepsilon.$$

It follows from (5.2) that there is $j_2 = j_2(\varepsilon, n, t, s) \in \mathbb{N}$ such that

$$|\nu_n(f_j, g_j; T_t^-) - \beta_n(t)| \le \varepsilon \quad \text{and} \quad |\nu_n(f_j, g_j; T_s^-) - \beta_n(s)| \le \varepsilon \quad \forall\, j \ge j_2.$$

Assuming that $s < t$ (the arguments are similar if $t < s$) and applying Lemma 2(a), (b), we get, for all $j \ge \max\{j_1, j_2\}$:

$$\begin{aligned}
|(f_j, g_j)(\{s,t\})| &\le \nu_{n+1}(f_j, g_j; T_t^-) - \nu_n(f_j, g_j; T_s^-) \\
&\le \nu_{n+1}(f_j, g_j; T_t^-) - \nu_n(f_j, g_j; T_t^-) \\
&\quad + |\nu_n(f_j, g_j; T_t^-) - \beta_n(t)| + |\beta_n(t) - \beta_n(s)| \\
&\quad + |\beta_n(s) - \nu_n(f_j, g_j; T_s^-)| \\
&\le \frac{\nu_{n+1}(f_j, g_j; T_t^-)}{n+1} + \varepsilon + \varepsilon + \varepsilon \\
&\le \frac{\nu_{n+1}(f_j, g_j)}{n+1} + 3\varepsilon \\
&\le \frac{\mu_{n+1}}{n+1} + \frac{\varepsilon}{n+1} + 3\varepsilon \le 5\varepsilon.
\end{aligned}$$

Being convergent (see condition (b)), sequences $\{f_j(s)\}_{j=1}^{\infty}$, $\{g_j(s)\}_{j=1}^{\infty}$ and $\{g_j(t)\}_{j=1}^{\infty}$ are Cauchy in $M$, and so, there is a number $j_3 = j_3(\varepsilon, s, t) \in \mathbb{N}$ such that, for all $j, k \ge j_3$, we have

$$d(f_j(s), f_k(s)) \le \varepsilon, \quad d(g_j(s), g_k(s)) \le \varepsilon, \quad \text{and} \quad d(g_j(t), g_k(t)) \le \varepsilon.$$

By virtue of inequality (2.3), we get

$$|(g_j, g_k)(\{s,t\})| \le d(g_j(s), g_k(s)) + d(g_j(t), g_k(t)) \le 2\varepsilon \quad \forall\, j, k \ge j_3.$$

Putting $j_4 = \max\{j_1, j_2, j_3\}$ and applying Lemma 1(e), (c), (b), we find

$$\begin{aligned}
d(f_j(t), f_k(t)) &\le d(f_j(s), f_k(s)) + |(f_j, f_k)(\{s,t\})| \\
&\le d(f_j(s), f_k(s)) + |(f_j, g_j)(\{s,t\})| + |(g_j, g_k)(\{s,t\})| \\
&\quad + |(g_k, f_k)(\{s,t\})| \\
&\le \varepsilon + 5\varepsilon + 2\varepsilon + 5\varepsilon = 13\varepsilon \quad \text{for all} \quad j, k \ge j_4.
\end{aligned}$$

Since $j_4$ depends only on $\varepsilon$ (and $t$), this proves that $\{f_j(t)\}_{j=1}^{\infty}$ is a Cauchy sequence in $M$, which together with assumption (a) establishes its convergence in $M$ to an element denoted by $f(t) \in M$.

Here and at the end of Step 3, we have shown that $f : T = S \cup (T \setminus S) \to M$ is a pointwise limit on $T$ of a subsequence $\{f_{j_k}\}_{k=1}^{\infty}$ of the original sequence $\{f_j\}_{j=1}^{\infty}$. Since $g_{j_k} \to g$ pointwise on $T$ as $k \to \infty$ as well, we conclude from Lemma 2(c) that

$$\nu_n(f,g) \le \liminf_{k\to\infty} \nu_n(f_{j_k}, g_{j_k}) \le \limsup_{j\to\infty} \nu_n(f_j, g_j) = \mu_n \quad \forall\, n \in \mathbb{N},$$

and so, $\nu_n(f,g) = o(n)$, or $f \in \mathrm{R}(g)$. This completes the proof of Theorem 1. $\square$

**Remark 3.** (a) Condition (b) in Theorem 1 may be replaced by the following one: $\{g_j(t)\}_{j=1}^{\infty}$ is a *Cauchy sequence* in $M$ for every $t \in T$. However, if $(M,d)$ is not complete, we may no longer infer the property $\nu_n(f,g) \le \mu_n$, $n \in \mathbb{N}$, of the pointwise limit $f$ (as there may be no $g$).

(b) Condition (2.6) is *necessary* for the uniformly convergent sequences $\{f_j\}$ and $\{g_j\}$: in fact, if $f_j \rightrightarrows f$ and $g_j \rightrightarrows g$ on $T$, and $\nu_n(f,g) = o(n)$, then it follows from Lemma 2(d) that $\lim_{j\to\infty} \nu_n(f_j, g_j) = \nu_n(f,g) = o(n)$.

(c) Condition (2.6) is 'almost necessary' in the following sense. Suppose $T \subset \mathbb{R}$ is a measurable set with Lebesgue measure $\mathcal{L}(T) < \infty$, $\{f_j\}$, $\{g_j\} \subset M^T$ are two sequences of measurable functions, which converge pointwise (or almost everywhere) on $T$ to functions $f, g \in M^T$, respectively, such that $\nu_n(f,g) = o(n)$. By Egorov's Theorem, given $\varepsilon > 0$, there exists a measurable set $E_\varepsilon \subset T$ such that $\mathcal{L}(T \setminus E_\varepsilon) \le \varepsilon$, $f_j \rightrightarrows f$ and $g_j \rightrightarrows g$ on $E_\varepsilon$. So, as in the previous remark (b), we have

$$\lim_{j\to\infty} \nu_n(f_j, g_j; E_\varepsilon) = \nu_n(f, g; E_\varepsilon) \le \nu_n(f,g) = o(n).$$

**Example 2.** (a) Condition (2.6) is *not necessary* for the pointwise convergence even if $g_j = c$ for all $j \in \mathbb{N}$. To see this, let $T = [0,1]$ and $x, y \in M$ with $x \ne y$. Given $j \in \mathbb{N}$, define $f_j : T \to M$ by: $f_j(t) = x$ if $j! \cdot t$ is integer, and $f_j(t) = y$ otherwise, $t \in [0,1]$. The pointwise precompact sequence $\{f_j\} \subset M^T$ consists of bounded regulated functions (in fact, $\nu_n(f_j, c) = o(n)$, and so, $f_j \in \mathrm{Reg}([0,1]; M) = \mathrm{R}(c)$ for all $j \in \mathbb{N}$). It converges pointwise on $T$ to the Dirichlet-type function $\mathcal{D}_{x,y}$ from (4.3). Note that $\nu_n(\mathcal{D}_{x,y}, c) = nd(x,y)$, and so, $\mathcal{D}_{x,y} \notin \mathrm{R}(c)$. Since the usual Jordan variation $V(f_j)$ of $f_j$ on $T = [0,1]$ is equal to $2 \cdot j! d(x,y)$, we find

$$\nu_n(f_j, c) = d(x,y) \cdot \begin{cases} n & \text{if} \quad n < 2 \cdot j!, \\ 2 \cdot j! & \text{if} \quad n \ge 2 \cdot j!, \end{cases} \qquad n, j \in \mathbb{N}.$$

Thus, $\lim_{j\to\infty} \nu_n(f_j, c) = d(x,y) \cdot n$, i.e., condition (2.6) does not hold.

(b) Under the assumptions of Theorem 1, condition (2.6) does not in general imply $\limsup_{j\to\infty}\nu_n(f_j,g)=o(n)$. To see this, let $g_j=f_j$ be as in example (a) above, so that $g=\mathcal{D}_{x,y}$. Given $n,j\in\mathbb{N}$, choose a collection $\{I_i\}_1^n\prec(0,1/j!)$ with $I_i=\{s_i,t_i\}$ such that $s_i$ is rational and $t_i$ is irrational for all $i=1,\dots,n$. Noting that, by virtue of (2.1),

$$|(f_j,g)(\{s_i,t_i\})|=\sup_{z\in M}|d(y,z)-d(x,z)|=d(y,x),$$

we get

$$\nu_n(f_j,g)\ge\sum_{i=1}^n|(f_j,g)(I_i)|=nd(y,x)\quad\text{for all}\quad n,j\in\mathbb{N}.$$

(c) The choice of an appropriate sequence $\{g_j\}$ is essential in Theorem 1. Let $\{x_j\}$, $\{y_j\}\subset M$ be two sequences, which converge in $M$ to $x,y\in M$, respectively, $x\ne y$. Define $f_j:T=[0,1]\to M$ by $f_j=\mathcal{D}_{x_j,y_j}$, $j\in\mathbb{N}$ (cf. (4.3)). Clearly, $\{f_j\}$ converges uniformly on $T$ to $\mathcal{D}_{x,y}$ (so, $\{f_j\}$ is pointwise precompact on $T$), and $\nu_n(f_j,c)=nd(x_j,y_j)$ for all $n,j\in\mathbb{N}$. Since

$$|d(x_j,y_j)-d(x,y)|\le d(x_j,x)+d(y_j,y)\to0\quad\text{as}\quad j\to\infty,$$

we find $\lim_{j\to\infty}\nu_n(f_j,c)=nd(x,y)$, condition (2.6) is not satisfied, and Theorem 1 is inapplicable with $g_j=c$, $j\in\mathbb{N}$.

On the other hand, set $g_j=\mathcal{D}_{x,y}$ for all $j\in\mathbb{N}$. Given $\{s,t\}\subset T$, we have, by virtue of (2.3),

$$|(f_j,g_j)(\{s,t\})|\le d(f_j(s),g_j(s))+d(f_j(t),g_j(t))\le2\varepsilon_j,$$

where $\varepsilon_j=\max\{d(x_j,x),d(y_j,y)\}\to0$ as $j\to\infty$. This implies the inequality $\nu_n(f_j,g_j)\le2n\varepsilon_j$, and so, condition (2.6) is fulfilled.

## 6. Extensions of known selection theorems

In this section, we consider extensions of two selection theorems from [25] and [22, 32]. The other selection theorems from the references in the Introduction were shown to be particular cases of [11]–[13] (see Remark 5).

By virtue of the inequality $\nu_n(f_j,g_j)/n\le V(f_j,g_j)/n$, instead of condition (2.6) in Theorem 1 we may assume that $\limsup_{j\to\infty}V(f_j,g_j)<\infty$ or $\sup_{j\in\mathbb{N}}V(f_j,g_j)<\infty$, in which cases the resulting pointwise limit $f$ of a subsequence of $\{f_j\}$ satisfies the regularity condition of the form $V(f,g)<\infty$.

Making use of the notion of $\varepsilon$-variation (Section 4), we get the following

**Theorem 3.** *Given $\varnothing \neq T \subset \mathbb{R}$ and a metric space $(M,d)$, let $\{f_j\} \subset M^T$ be a pointwise precompact sequence of functions such that*

$$\limsup_{j\to\infty} V_\varepsilon(f_j) < \infty \quad \textit{for all} \quad \varepsilon > 0. \tag{6.1}$$

*Then, there is a subsequence $\{f_{j_k}\}$ of $\{f_j\}$, which converges pointwise on $T$ to a regulated function $f \in \mathrm{R}(c)$.*

**Proof**. Taking into account Theorem 1, it suffices to verify that condition (6.1) implies $\limsup_{j\to\infty} \nu_n(f_j,c) = o(n)$, which is (2.6) with $g_j = c$ for all $j \in \mathbb{N}$. In fact, by (6.1), for every $\varepsilon > 0$ there is $j_0 = j_0(\varepsilon) \in \mathbb{N}$ and $C(\varepsilon) > 0$ such that $V_\varepsilon(f_j) \le C(\varepsilon)$ for all $j \ge j_0$. Definition (4.1) yields the existence of $g_j \in \mathrm{BV}(T;M)$ such that $d_\infty(f_j,g_j) \le \varepsilon$ and $V(g_j) \le V_\varepsilon(f_j) + (1/j) \le C(\varepsilon)+1$ for all $j \ge j_0$. By (3.7) (where we replace $g_j$ and $g$ by $c$, and $f$—by $g_j$),

$$\frac{\nu_n(f_j,c)}{n} \le 2d_\infty(f_j,g_j) + \frac{\nu_n(g_j,c)}{n} \le 2\varepsilon + \frac{V(g_j)}{n} \le 2\varepsilon + \frac{C(\varepsilon)+1}{n}$$

for all $j \ge j_0$ and $n \in \mathbb{N}$. Consequently,

$$\frac{1}{n}\limsup_{j\to\infty} \nu_n(f_j,c) \le \frac{1}{n}\sup_{j\ge j_0} \nu_n(f_j,c) \le 2\varepsilon + \frac{C(\varepsilon)+1}{n} \quad \forall\, \varepsilon > 0,\ n \in \mathbb{N}.$$

This implies that the left-hand side in this inequality tends to zero as $n \to \infty$: given $\eta > 0$, we set $\varepsilon = \eta/4$ and choose a number $n_0 = n_0(\eta) \in \mathbb{N}$ such that $(C(\varepsilon)+1)/n_0 \le \eta/2$, which yields $2\varepsilon + (C(\varepsilon)+1)/n \le \eta$ for all $n \ge n_0$. □

**Remark 4.** (a) If $M = \mathbb{R}^N$ in Theorem 3, we may infer that $V_\varepsilon(f)$ does not exceed the limit superior from (6.1): in fact, it follows from [25, Proposition 3.6] that $V_\varepsilon(f) \le \liminf_{k\to\infty} V_\varepsilon(f_{j_k})$ for all $\varepsilon > 0$.

(b) Theorem 3 extends Theorem 3.8 from [25], which has been established for $T = [a,b]$ and $M = \mathbb{R}^N$ under the assumption that $\sup_{j\in\mathbb{N}} V_\varepsilon(f_j) < \infty$ for every $\varepsilon > 0$. The last assumption on the uniform boundedness of $\varepsilon$-variations is more restrictive than condition (6.1) as the following example shows.

**Example 3.** Let $\{x_j\}$ and $\{y_j\}$ be two sequences from $M$ such that $x_j \neq y_j$ for all $j \in M$ and, for some $x \in M$, $x_j \to x$ and $y_j \to x$ in $M$ as $j \to \infty$.

We set $f_j = \mathcal{D}_{x_j,y_j}$, $j \in \mathbb{N}$, and $f(t) = x$ for all $t \in T = [0,1]$. The sequence $\{f_j\} \subset M^T$ converges uniformly on $T$ to the constant function $f$:

$$d_\infty(f_j, f) = \max\{d(x_j, x), d(y_j, x)\} \to 0 \quad \text{as} \quad j \to \infty.$$

Given $\varepsilon > 0$, there is $j_0 = j_0(\varepsilon) \in \mathbb{N}$ such that $d(x_j, y_j) \le \varepsilon$ for all $j \ge j_0$, and so, by (4.4), $V_\varepsilon(f_j) = 0$ for all $j \ge j_0$, which implies condition (6.1):

$$\limsup_{j\to\infty} V_\varepsilon(f_j) \le \sup_{j\ge j_0} V_\varepsilon(f_j) = 0.$$

On the other hand, if $k \in \mathbb{N}$ is fixed and $0 < \varepsilon < d(x_k, y_k)/2$, then (4.4) gives $V_\varepsilon(f_k) = \infty$, and so, $\sup_{j\in\mathbb{N}} V_\varepsilon(f_j) = \infty$.

Now, we are going to present an extension of a Helly-type selection theorem from [32, Section 4, Theorem 1] and [22, Theorem 2].

Let $\kappa : [0,1] \to [0,1]$ be a continuous, increasing and concave function such that $\kappa(0) = 0$, $\kappa(1) = 1$, and $\kappa(\tau)/\tau \to \infty$ as $\tau \to +0$ (e.g., $\kappa(\tau) = \tau(1 - \log\tau)$, $\kappa(\tau) = \tau^\alpha$ with $0 < \alpha < 1$, or $\kappa(\tau) = 1/(1 - \frac{1}{2}\log\tau)$, see [33]).

Let $T = [a,b]$ be a closed interval in $\mathbb{R}$, $a < b$. We set $|T| = b - a$, and if $\{t_i\}_0^n \prec [a,b]$ is a partition of $T$ (i.e., $a = t_0 < t_1 < \cdots < t_{n-1} < t_n = b$), we also set $I_i = \{t_{i-1}, t_i\}$ and $|I_i| = t_i - t_{i-1}$, $i = 1, \ldots, n$.

The *joint $\kappa$-variation* of functions $f, g \in M^T = M^{[a,b]}$ is defined by

$$V_\kappa(f,g) = \sup\left\{ \sum_{i=1}^n |(f,g)(I_i)| \Big/ \sum_{i=1}^n \kappa\big(|I_i|/|T|\big) : n \in \mathbb{N} \text{ and } \{t_i\}_0^n \prec [a,b] \right\},$$

where $|(f,g)(I_i)| = |(f,g)(\{t_{i-1}, t_i\})|$ is given by (2.1).

Since $|(f,c)(I_i)| = d(f(t_{i-1}), f(t_i))$ is independent of a constant function $c : [a,b] \to M$, the quantity $V_\kappa(f) \equiv V_\kappa(f,c)$ is the *Korenblum $\kappa$-variation* of $f \in M^{[a,b]}$, introduced in [32, p. 191] and [33, Section 5] for $M = \mathbb{R}$.

The following theorem is a generalization of Theorem 2 from [22], established for real functions of bounded $\kappa$-variation under the assumption that $\sup_{j\in\mathbb{N}} V_\kappa(f_j) < \infty$ and based on the decomposition of any $f \in \mathbb{R}^{[a,b]}$ with $V_\kappa(f) < \infty$ into the difference of two real $\kappa$-decreasing functions.

**Theorem 4.** *Under the assumptions of Theorem 1, suppose that condition* (2.6) *is replaced by the following one*: $\limsup_{j\to\infty} V_\kappa(f_j, g_j) < \infty$. *Then, there is a subsequence of $\{f_j\}$, which converges pointwise on $T = [a,b]$ to a function $f \in \mathrm{R}(g)$ such that $V_\kappa(f,g) < \infty$.*

**Proof**. In order to show that (2.6) is satisfied, let $n \in \mathbb{N}$, $\{I_i\}_1^n \prec [a,b]$ with $I_i = \{s_i, t_i\}$, and set $I_i' = \{t_i, s_{i+1}\}$ and $|I_i'| = s_{i+1} - t_i$, $i = 1, \dots, n-1$. By the definition of $V_\kappa(f_j, g_j)$ and the concavity of function $\kappa$, we have

$$\begin{aligned}
\sum_{i=1}^{n} |(f_j, g_j)(I_i)| \leq & |(f_j, g_j)(\{a, s_1\})| + \sum_{i=1}^{n} |(f_j, g_j)(I_i)| \\
& + \sum_{i=1}^{n-1} |(f_j, g_j)(I_i')| + |(f_j, g_j)(\{t_n, b\})| \\
& \leq \left[ \kappa\left(\frac{s_1 - a}{|T|}\right) + \sum_{i=1}^{n} \kappa\left(\frac{|I_i|}{|T|}\right) + \sum_{i=1}^{n-1} \kappa\left(\frac{|I_i'|}{|T|}\right) + \kappa\left(\frac{b - t_n}{|T|}\right) \right] V_\kappa(f_j, g_j) \\
& \leq (2n+1)\kappa\left( \frac{1}{(2n+1)(b-a)} \left[ (s_1 - a) + \sum_{i=1}^{n} (t_i - s_i) \right.\right. \\
& \quad \left.\left. + \sum_{i=1}^{n-1} (s_{i+1} - t_i) + (b - t_n) \right] \right) V_\kappa(f_j, g_j) \\
& \leq (2n+1)\kappa\left( \frac{1}{2n+1} \right) V_\kappa(f_j, g_j).
\end{aligned}$$

Thus,

$$\frac{\nu_n(f_j, g_j)}{n} \leq \left(2 + \frac{1}{n}\right) \kappa\left(\frac{1}{2n+1}\right) V_\kappa(f_j, g_j) \quad \text{for all} \quad j, n \in \mathbb{N},$$

and so, condition (2.6) in Theorem 1 is satisfied.

Let $f \in \mathrm{R}(g)$ be the pointwise limit of a subsequence $\{f_{j_m}\}$ of $\{f_j\}$. Arguing as in the proof of Lemma 2(c), we get

$$V_\kappa(f, g) \leq \liminf_{m \to \infty} V_\kappa(f_{j_m}, g_{j_m}) \leq \limsup_{j \to \infty} V_\kappa(f_j, g_j) < \infty. \qquad \square$$

**Remark 5.** Since Theorem 1 is an extension of results from [11, 12], it also contains as particular cases all pointwise selection principles based on various notions of generalized variation. These principles may be further generalized in the spirit of Theorem 4 replacing the increment $|f(I_i)| = d(f(s_i), f(t_i))$ applied in [11]–[13] by the joint increment $|(f, g)(I_i)|$ from (2.1).

Finally, it is worth mentioning that the joint modulus of variation (2.4), defined by means of (2.1), plays an important role in the extension of a result from [17] to metric space valued functions:

**Theorem 5.** *Given $\varnothing \neq T \subset \mathbb{R}$ and a metric space $(M, d)$, let $\{f_j\} \subset M^T$ be a pointwise precompact sequence of functions such that*

$$\lim_{N\to\infty} \sup_{j,k\geq N} \nu_n(f_j, f_k) = o(n).$$

*Then, there is a subsequence of $\{f_j\}$, which converges pointwise on $T$.*

Taking into account Lemmas 1 and 2, the proof of this theorem follows the same lines as the ones given in [17, Theorem 1] (where $(M, d, +)$ is a metric semigroup and $\nu_n(f_j, f_k)$ is defined via (3.1)), and so, it is omitted. Note that the limit of a pointwise convergent subsequence of $\{f_j\}$ in Theorem 5 may be a non-regulated function. For more details, examples and relations with previously known 'regular' and 'irregular' versions of pointwise selection principles from [23, 40, 42] we refer to [14, Section 5.2], [17, 18, 34].

## 7. Conclusions

In the context of functions of one real variable taking values in a metric space, we have presented a pointwise selection principle, which can be applied to arbitrary (regulated and non-regulated) sequences of functions. It is based on notions of joint increment and joint modulus of variation, the latter being a nondecreasing sequence of pseudometrics on the appropriate product space. Our selection principle extends the classical Helly Theorem and contains as particular cases many selection theorems based on various notions of generalized variation of functions. In contrast to previously established selection principles, the main assumption in our theorem on the limit superior is 'almost necessary', and it is shown by examples that in a certain sense this assumption is sharp. The notion of conditionally regulated functions explains the limitations of previously known selection results.

## References


[1] V. Barbu, Th. Precupanu, Convexity and Optimization in Banach Spaces, second ed., Reidel, Dordrecht, 1986.

[2] S. A. Belov, V. V. Chistyakov, A selection principle for mappings of bounded variation, J. Math. Anal. Appl. 249 (2) (2000) 351–366; and comments on this paper: J. Math. Anal. Appl. 278 (1) (2003) 250–251.

[3] P. C. Bhakta, On functions of bounded variation relative to a set, J. Austral. Math. Soc. 13 (1972) 313–322.

[4] Z. A. Chanturiya, The modulus of variation of a function and its application in the theory of Fourier series, Soviet Math. Dokl. 15 (1974) 67–71.

[5] Z. A. Chanturiya, Absolute convergence of Fourier series, Math. Notes 18 (1975) 695–700.

[6] V. V. Chistyakov, On mappings of bounded variation, J. Dynam. Control Systems 3 (2) (1997) 261–289.

[7] V. V. Chistyakov, On the theory of multivalued mappings of bounded variation of one real variable, Sb. Math. 189 (5–6) (1998) 797–819.

[8] V. V. Chistyakov, Mappings of bounded variation with values in a metric space: generalizations, in: Pontryagin Conference, 2, Nonsmooth Analysis and Optimization (Moscow, 1998). J. Math. Sci. (New York) 100 (6) (2000) 2700–2715.

[9] V. V. Chistyakov, On multi-valued mappings of finite generalized variation, Math. Notes 71 (3–4) (2002) 556–575.

[10] V. V. Chistyakov, Selections of bounded variation, J. Appl. Anal. 10 (1) (2004) 1–82.

[11] V. V. Chistyakov, The optimal form of selection principles for functions of a real variable, J. Math. Anal. Appl. 310 (2) (2005) 609–625.

[12] V. V. Chistyakov, A selection principle for functions of a real variable, Atti Sem. Mat. Fis. Univ. Modena e Reggio Emilia 53 (1) (2005) 25–43.

[13] V. V. Chistyakov, A pointwise selection principle for functions of a real variable with values in a uniform space, Siberian Adv. Math. 16 (3) (2006) 15–41.

[14] V. V. Chistyakov, Metric Modular Spaces: Theory and Applications, SpringerBriefs in Mathematics, Springer, Switzerland, 2015.

[15] V. V. Chistyakov, O. E. Galkin, On maps of bounded $p$-variation with $p > 1$, Positivity 2 (1) (1998) 19–45.

[16] V. V. Chistyakov, O. E. Galkin, Mappings of bounded $\Phi$-variation with arbitrary function $\Phi$, J. Dynam. Control Systems 4 (2) (1998) 217–247.

[17] V. V. Chistyakov, C. Maniscalco, A pointwise selection principle for metric semigroup valued functions, J. Math. Anal. Appl. 341 (1) (2008) 613–625.

[18] V. V. Chistyakov, C. Maniscalco, Yu. V. Tretyachenko, Variants of a selection principle for sequences of regulated and non-regulated functions, in: L. De Carli, K. Kazarian and M. Milman (Eds.), Topics in Classical Analysis and Applications in Honor of Daniel Waterman, World Scientific Publishing, Hackensack, NJ, 2008, pp. 45–72.

[19] V. V. Chistyakov, A. Nowak, Regular Carathéodory-type selectors under no convexity assumptions, J. Funct. Anal. 225 (2) (2005) 247–262.

[20] V. V. Chistyakov, D. Repovš, Selections of bounded variation under the excess restrictions, J. Math. Anal. Appl. 331 (2) (2007) 873–885.

[21] V. V. Chistyakov, A. Rychlewicz, On the extension and generation of set-valued mappings of bounded variation, Studia Math. 153 (3) (2002) 235–247.

[22] D. S. Cyphert, J. A. Kelingos, The decomposition of functions of bounded $\varkappa$-variation into differences of $\varkappa$-decreasing functions, Studia Math. 81 (2) (1985) 185–195.

[23] L. Di Piazza, C. Maniscalco, Selection theorems, based on generalized variation and oscillation, Rend. Circ. Mat. Palermo, Ser. II, 35 (3) (1986) 386–396.

[24] R. Duda, On convex metric spaces. III, Fund. Math. 51 (1962) 23–33.

[25] D. Fraňková, Regulated functions, Math. Bohem. 116 (1) (1991) 20–59.

[26] S. Gniłka, On the generalized Helly's theorem, Funct. Approx. Comment. Math. 4 (1976) 109–112.

[27] K. Goebel, W. A. Kirk, Topics in Metric Fixed Point Theory, Cambridge Stud. Adv. Math., vol. 28, Cambridge Univ. Press, Cambridge, 1990.

[28] C. Goffman, T. Nishiura, D. Waterman, Homeomorphisms in Analysis, Math. Surveys Monogr., vol. 54, Amer. Math. Soc., Providence, Rhode Island, 1997.

[29] E. Helly, Über lineare Funktionaloperationen, Sitzungsber. Naturwiss. Kl. Kaiserlichen Akad. Wiss. Wien 121 (1912) 265–297.

[30] H. Hermes, On continuous and measurable selections and the existence of solutions of generalized differential equations, Proc. Amer. Math. Soc. 29 (3) (1971) 535–542.

[31] T. H. Hildebrandt, Introduction to the Theory of Integration, Academic Press, New York, London, 1963.

[32] B. Korenblum, An extension of the Nevanlinna theory, Acta Math. 135 (1) (1975) 187–219.

[33] B. Korenblum, A generalization of two classical convergence tests for Fourier series, and some new Banach spaces of functions, Bull. Amer. Math. Soc. (N. S.) 9 (2) (1983) 215–218.

[34] C. Maniscalco, A comparison of three recent selection theorems, Math. Bohem. 132 (2) (2007) 177–183.

[35] K. Menger, Untersuchungen über allgemeine Metrik, Math. Ann. 100 (1928) 75–163.

[36] B. S. Mordukhovich, Approximation Methods in the Problems of Optimization and Control, Nauka, Moscow, 1988 (in Russian).

[37] P. Muldowney, A Modern Theory of Random Variation: With Applications in Stochastic Calculus, Financial Mathematics, and Feynman Integration, John Wiley & Sons, Hoboken, New Jersey, 2012.

[38] J. Musielak, W. Orlicz, On generalized variations (I), Studia Math. 18 (1959) 11–41.

[39] I. P. Natanson, Theory of Functions of a Real Variable, third ed., Nauka, Moscow, 1974 (in Russian). English translation: Ungar, New York, 1965.

[40] K. Schrader, A generalization of the Helly selection theorem, Bull. Amer. Math. Soc. 78 (3) (1972) 415–419.

[41] M. Schramm, Functions of $\Phi$-bounded variation and Riemann-Stieltjes integration, Trans. Amer. Math. Soc. 287 (1985) 49–63.

[42] Yu. V. Tret'yachenko, V. V. Chistyakov, The selection principle for pointwise bounded sequences of functions, Math. Notes 84 (4) (2008) 396–406.

[43] D. Waterman, On $\Lambda$-bounded variation, Studia Math. 57 (1976) 33–45.